

Spin dependent Fermi Liquid parameters and properties of polarized quark matter

Kausik Pal,^{*} Subhrajyoti Biswas, and Abhee K. Dutt-Mazumder

*High Energy Physics Division, Saha Institute of Nuclear Physics,
1/AF Bidhannagar, Kolkata 700064, India.*

We calculate the spin dependent Fermi liquid parameters (FLPs), single particle energies and energy densities of various spin states of polarized quark matter. The expressions for the incompressibility (K) and sound velocity (c_1) in terms of the spin dependent FLPs and polarization parameter (ξ) are derived. Estimated values of K and c_1 reveal that the equation of state (EOS) of the polarized matter is stiffer than the unpolarized one. Finally we investigate the possibility of the spin polarization (ferromagnetism) phase transition.

PACS numbers: 04.40.Dg, 12.38.Bx, 12.39.-x, 14.70.Dj, 26.60.-c, 97.60.Jd

Keywords: Ferromagnetism, Quark matter, Landau parameters.

I. INTRODUCTION

One of the important research areas of the contemporary high energy physics has been the study of matter under extreme conditions. Such a matter, in the laboratory can be produced by colliding heavy ions at ultra-relativistic energies. Due to asymptotic freedom of quantum chromodynamics (QCD), it is predicted that the hadronic matter at high temperature and/or density can undergo a series of phase transitions like confinement-deconfinement and/or chiral phase transition [1, 2]. In the high density regime QCD predicts the existence of color superconducting state [3, 4, 5]. These apart, the possibility of spin polarized quark liquid *i.e.* the existence of ferromagnetic phase in dense quark system has also been suggested recently with which we are presently concerned [2, 6]. The properties of dense quark system are particularly relevant for the study of various astrophysical phenomenon.

^{*}Electronic address: kausik.pal@saha.ac.in

The part of the motivation to study the ferromagnetic phase transition in dense quark matter (DQM), as mentioned in [6] is provided by the discovery of ‘magnetars’ [7] where an extraordinarily high magnetic field $\sim 10^{15}G$ exists [6, 8]. In [6], it is argued, that the origin of such a high magnetic field can be attributed to the existence of spin polarized quark matter [9]. To examine the possibility of ferromagnetism in DQM in ref.[6] a variational calculation is performed where it is observed that there exists a critical density below which spin polarized quark matter is energetically favorable than unpolarized state. Subsequently various other calculations were also performed to investigate this issue [2, 4, 5, 7, 8, 9, 10]. For example, in [5] it is shown that there is no contradiction between color superconductivity and ferromagnetism and both of these phase can co-exist. In [10], the same problem was studied in the large N_c and N_f limit while keeping N_c/N_f fixed where it was shown that spin polarized state can exist, however, in presence of magnetic screening, color superconductivity or dense chiral waves disappear. It might be mentioned that such screening is now supported by the lattice calculation [10, 11]. In[5] it is analytically shown that, if quarks are massless, ferromagnetism does not appear which is consistent with the conclusion drawn in [10]. Ref.[8] shows that ferromagnetism might appear in quark matter with Goldstone boson current where the magnetization is shown to be related to triangle anomalies.

In the present work, we apply relativistic Fermi liquid theory (RFLT) to study the possibility of para-ferro phase transition in DQM. The relativistic Fermi liquid theory was developed by Baym and Chin [14] where it has been shown how the various physical quantities like chemical potential (μ), incompressibility (K), sound velocity (c_1) etc. can be expressed in terms of the Landau parameters (LPs) calculated relativistically. However, the formalism developed in [14] is valid for unpolarized matter and LPs calculated there are spin averaged.

In this paper we extend the formalism of RFLT and the required LPs are calculated by retaining their explicit spin dependencies. As a result, here various combination of parameters like $f_{0,1}^{++}$, $f_{0,1}^{+-}$, $f_{0,1}^{-+}$ and $f_{0,1}^{--}$ corresponding to scattering involving up-up, up-down, down-up or down-down spins are appear [14]. Once determined, these parameters are used to calculate quantities like chemical potentials for the spin up and spin down quarks or the total energy density of the system as a function of $\xi = (n_q^+ - n_q^-)/n_q$ and n_q together with various other quantities as we shall see. Here n_q^+ and n_q^- correspond to densities of spin up, down quarks respectively and $n_q = n_q^+ + n_q^-$, denotes total quark density [6]. We, also compare some of our results with those presented in [6] where more direct approach was

adopted to calculate the total energy density from the loop. In addition, the present work is extended further to estimate incompressibility and sound velocity in dense quark system for a given fraction of spin-up or down quarks.

Furthermore, in dealing with the massless gluons, we find that naive series expansion fails and one has to use hard density loop (HDL) corrected gluon propagator to get the finite result for the LPs involving scattering of like spins [12]. This however does not cause any problem for the calculation of various physical quantities like chemical potential, exchange energy, incompressibility etc. We shall see, even though f_0 and f_1 (suppressing spin indices) individually remain divergent, what appears in our case is the particular combination of these parameters where such divergences cancel.

The plan of the paper is as follows. In Sec.II, as mentioned before, we extend the formalism of RFLT to include explicit spin dependence. In Sec.III, we derive spin dependent LPs due to one gluon exchange (OGE) for polarized quark matter. Subsequently, we calculate chemical potential and energy density. We find the density dependence of incompressibility (K) and first sound velocity (c_1) with arbitrary spin polarization (ξ). To compare with ref.[6], we present ultra-relativistic and non-relativistic results and studied para-ferro phase transition of quark matter. Sec.IV is devoted to summary and conclusion. In Appendix, we calculate various LPs for unlike spin states of scatterer.

II. FORMALISM

In FLT total energy density E of an interacting system is the functional of occupation number n_p of the quasi-particle states of momentum p . The excitation of the system is equivalent to the change of occupation number by an amount δn_p . The corresponding energy density of the system is given by [13, 14],

$$E = E^0 + \sum_s \int \frac{d^3p}{(2\pi)^3} \varepsilon_{ps}^0 \delta n_{ps} + \frac{1}{2} \sum_{ss'} \int \frac{d^3p}{(2\pi)^3} \frac{d^3p'}{(2\pi)^3} f_{ps,p's'} \delta n_{ps} \delta n_{p's'}, \quad (1)$$

where E^0 is the ground state energy density and s is the spin index, and the quasi-particle energy can be written as,

$$\varepsilon_{ps} = \varepsilon_{ps}^0 + \sum_{s'} \int \frac{d^3p'}{(2\pi)^3} f_{ps,p's'} \delta n_{p's'}, \quad (2)$$

where ε_{ps}^0 is the non-interacting single particle energy. The interaction between quasiparticles is given by $f_{ps,p's'}$, which is defined to be the second derivative of the energy functional with respect to occupation functions,

$$f_{ps,p's'} = \frac{\delta^2 E}{\delta n_{ps} \delta n_{p's'}}. \quad (3)$$

Since, the quasiparticles are well defined only near the Fermi surface, one assumes

$$\varepsilon_{ps} = \mu^s + v_f^s(p - p_f^s). \quad (4)$$

In FLT, the interaction parameter, $f_{ps,p's'}$, is expanded on the basis of Legendre polynomials, P_l [13, 14]. The coefficients of this expansion are known as FLPs, which are given by

$$f_l^{ss'} = (2l + 1) \int \frac{d\Omega}{4\pi} P_l(\cos \theta) f_{ps,p's'}, \quad (5)$$

where θ is the angle between p and p' , both taken to be on the Fermi surface, and the integration is over all directions of p [14]. Note that unlike [13, 14], here we retain explicit spin indices without performing spin summation. We restrict ourselves for $l \leq 1$ i.e. f_0^s and f_1^s , since higher l contribution decreases rapidly as the scattering is dominated by the small angles and the series converges, here, $f_l^s = \frac{1}{2} \sum_{s'} f_l^{ss'}$ [15].

The Landau Fermi liquid interaction $f_{ps,p's'}$ is related to the two particle forward scattering amplitude via [13, 14],

$$f_{ps,p's'} = \frac{m_q}{\varepsilon_p^0} \frac{m_q}{\varepsilon_{p'}^0} \mathcal{M}_{ps,p's'}, \quad (6)$$

where m_q is the mass of the quark and the Lorentz invariant matrix $\mathcal{M}_{ps,p's'}$ consists of the usual direct and exchange amplitude, which may, therefore be evaluated by conventional Feynman rules. The dimensionless LPs are defined as $F_l^s = N^s(0) f_l^s$ [14], where $N^s(0)$ is the density of states at the Fermi surface is given by,

$$\begin{aligned} N^s(0) &= \int \frac{d^3p}{(2\pi)^3} \delta(\varepsilon_{ps} - \mu^s) \\ &= \frac{g_{deg} p_f^{s^2}}{2\pi^2} \left(\frac{\partial p}{\partial \varepsilon_{ps}} \right)_{p=p_f^s} \\ &\simeq \frac{g_{deg} p_f^s \varepsilon_f^s}{2\pi^2}. \end{aligned} \quad (7)$$

Here g_{deg} is the degeneracy factor. In our case $g_{deg} = N_c N_f$ where N_c and N_f are the color and flavor index for quark matter. For spin up (+) and spin down (-) quark, density of states will be change accordingly. In the above expression $(\partial p / \partial \varepsilon_{ps})_{p=p_f^s}$ is the inverse Fermi velocity $(1/v_f^s)$ related to the FL parameter F_1^s ,

$$\frac{1}{v_f^s} = (\partial p / \partial \varepsilon_{ps})_{p=p_f^s} = (\mu^s / p_f^s)(1 + F_1^s/3). \quad (8)$$

With Eq.(7) and Eq.(8) one reads the general relation as [16]

$$\varepsilon_f^s = \mu^s(1 + \frac{1}{3}F_1^s). \quad (9)$$

The compression modulus or incompressibility (K) of the system is defined by the second derivative of total energy density E with respect to the number density n_q , is given by [16, 17, 18, 19, 20]

$$K = 9n_q \frac{\partial^2 E}{\partial n_q^2}. \quad (10)$$

Now we introduce a polarization parameter ξ by the equations, $n_q^+ = n_q(1 + \xi)/2$ and $n_q^- = n_q(1 - \xi)/2$ under the condition $0 \leq \xi \leq 1$ [6]. The Fermi momenta in the spin-polarized quark matter then are $p_f^+ = p_f(1 + \xi)^{1/3}$ and $p_f^- = p_f(1 - \xi)^{1/3}$, where $p_f = (\pi^2 n_q)^{1/3}$, is the Fermi momentum of the unpolarized matter ($\xi = 0$). So, there are two Fermi surfaces corresponding to spin-up (+) and spin-down (-) states, such that $E \equiv E(n_q^+, n_q^-)$. We have

$$\begin{aligned} \frac{\partial E}{\partial n_q} &= \frac{\partial E}{\partial n_q^+} \frac{\partial n_q^+}{\partial n_q} + \frac{\partial E}{\partial n_q^-} \frac{\partial n_q^-}{\partial n_q} \\ &= \frac{1}{2} [(1 + \xi)\mu^+ + (1 - \xi)\mu^-] \end{aligned} \quad (11)$$

Using Eq.(11), the incompressibility becomes[20]

$$\begin{aligned} K &= \frac{9n_q}{4} \left[(1 + \xi)^2 \frac{\partial \mu^+}{\partial n_q^+} + (1 - \xi)^2 \frac{\partial \mu^-}{\partial n_q^-} \right] \\ &= \frac{9n_q}{4} \left[(1 + \xi)^2 \left(\frac{1 + F_0^+}{N^+(0)} \right) + (1 - \xi)^2 \left(\frac{1 + F_0^-}{N^-(0)} \right) \right], \end{aligned} \quad (12)$$

where [14]

$$\frac{\partial \mu^s}{\partial n_q^s} = \frac{1 + F_0^s}{N^s(0)}. \quad (13)$$

Similarly, the relativistic first sound velocity is given by the first derivative of pressure P with respect to energy density E . Since $P = \sum_s \mu^s n_q^s - E$ [16, 20], we have,

$$\begin{aligned} c_1^2 &= \frac{\partial P}{\partial E} = \frac{\partial P}{\partial n_q} \frac{\partial n_q}{\partial E} \\ &= \left[\frac{(1 + \xi) n_q^+ \frac{\partial \mu^+}{\partial n_q^+} + (1 - \xi) n_q^- \frac{\partial \mu^-}{\partial n_q^-}}{(1 + \xi) \mu^+ + (1 - \xi) \mu^-} \right] \\ &= \frac{n_q}{2[(1 + \xi) \mu^+ + (1 - \xi) \mu^-]} \left[(1 + \xi)^2 \left(\frac{1 + F_0^+}{N^+(0)} \right) + (1 - \xi)^2 \left(\frac{1 + F_0^-}{N^-(0)} \right) \right]. \end{aligned} \quad (14)$$

In the above Eq.(12) and Eq.(14), $N^\pm(0)$ and $F_0^\pm$ corresponds to density of states at Fermi surface and dimensionless LP for spin up (+) and spin down (-) quark respectively. For unpolarized matter, $\xi = 0$ implying $\mu^+ = \mu^-$, $F_0^+ = F_0^-$ and $N^+(0) = N^-(0)$. From Eq.(12) and (14) we have the well known result as $K = 9n_q \frac{\partial \mu}{\partial n_q}$ [16] and $c_1^2 = \frac{n_q}{\mu} \frac{\partial \mu}{\partial n_q}$ [14].

III. LANDAU PARAMETERS FOR POLARIZED QUARK MATTER

In this section we calculate LPs for quark matter with explicit spin dependencies. We choose spin s along z axis *i.e.* $s \equiv (0, 0, \pm 1)$ and represent spin-up and down states by their signs. For a four-dimensional description of the polarization state, it is convenient to define a 4-vector a^μ which, in the rest frame of each quark, is same as the three-dimensional vector s ; since s is an axial vector, a^μ is a 4-pseudovector. This 4-vector is orthogonal to the 4-momentum in the rest frame (in which $a^\mu = (0, s)$, $P^\mu = (m_q, 0)$); in any frame we therefore have $a^\mu P_\mu = 0$ [6, 21, 22].

The components of the 4-vector a^μ in a frame in which the particle is moving with momentum p are found by a Lorentz transformation from the rest frame [22],

$$a = s + \frac{p(s \cdot p)}{m_q(\varepsilon_p + m_q)}; \quad a^0 = \frac{p \cdot s}{m_q} \quad (15)$$

with $\varepsilon_p = \sqrt{p^2 + m_q^2}$. We can define projection operator $P(a)$ on each of spin polarization, $P(a) = \frac{1}{2}(1 + \gamma^5 \not{a})$. Accordingly the polarization density matrix ρ is given by the expression

$$\rho(P, s) = \frac{1}{2m_q}(\not{P} + m_q)P(a), \quad (16)$$

which is normalized by the condition, $\text{Tr}\rho(P, s) = 1$. The mean value of the spin is then given by the quantity [22]

$$\begin{aligned} s_{av} &= \frac{1}{2} \frac{m_q}{\varepsilon_p} \text{Tr}(\rho \gamma_0 \Sigma) = \frac{1}{2} \frac{m_q}{\varepsilon_p} \text{Tr}(\rho \gamma_5 \gamma) \\ &= \frac{1}{2} \frac{m_q}{\varepsilon_p} \left(s + \frac{p(s \cdot p)}{m_q(\varepsilon_p + m_q)} \right), \end{aligned} \quad (17)$$

which is reduced to $s_{av} = \frac{1}{2}s$ in the non-relativistic limit.

We consider the color-symmetric forward scattering amplitude of the two quarks around the Fermi surface by the OGE interaction. The direct term does not contribute as it involves trace of single color matrices like $\text{Tr}\lambda_a$, which vanishes. Thus the leading contribution comes from the exchange (Fock) term [6]:

$$\begin{aligned} \mathcal{M}_{ps,p's'}^{ex} &= -\frac{1}{3} \sum_i \frac{1}{3} \sum_j [\bar{U}_\beta(P') g(t^a)_{ji} \gamma^\mu U_\alpha(P)] \left(\frac{-g_{\mu\nu}}{(P - P')^2} \right) [\bar{U}_\alpha(P) g(t^a)_{ij} \gamma^\nu U_\beta(P')] \\ &= \frac{4}{9} \frac{1}{(P - P')^2} \text{Tr}[\gamma_\mu \rho(P, s) \gamma^\mu \rho(P', s')], \end{aligned} \quad (18)$$

where α, β is the flavor level, i, j is the quark color index, $t^a (= \lambda_a/2)$ is the color matrix and g is the coupling constant. Since gluon is flavor blind, the u -channel diagrams contribute only when $\alpha = \beta$; *i.e.* scattering of quarks with same flavor [23]. This means that the Fermi sphere of each flavor makes an independent contribution. Thus the potential energy receives a factor N_f . On the other hand, the quarks with different colors can take part in the exchange process, giving rise to a factor N_c^2 . Eventually the potential energy density is proportional to $N_f N_c^2 g^2$. For the kinetic energy density, there arises an overall factor $N_c N_f$. Thus, the factor $N_c N_f$ factorizes out of the total energy density and the competition between the kinetic and potential energies is not influenced by the number of flavor. The number of flavor neither encourages or discourages ferromagnetism [10].

Without loss of generality, for the calculation of energy density and other related quantities, we consider one-flavor quark matter. With the help of polarization density matrices

given in Eq.(16), we have from Eq.(18) the interaction amplitude as [6]

$$\begin{aligned} \mathcal{M}_{ps,p's'}^{ex} = & \frac{2g^2}{9m_q^2} \frac{1}{(P-P')^2} [2m_q^2 - P \cdot P' - (p \cdot s)(p' \cdot s') + m_q^2(s \cdot s') + \frac{1}{(\varepsilon_p + m_q)(\varepsilon_{p'} + m_q)} \\ & \times \{m_q(\varepsilon_p + m_q)(p' \cdot s)(p' \cdot s') + m_q(\varepsilon_{p'} + m_q)(p \cdot s)(p \cdot s') + (p \cdot p')(p \cdot s)(p' \cdot s')\}. \end{aligned} \quad (19)$$

From Eq.(6) the quasiparticle interaction parameter is given by

$$f_{ps,p's'}^{ex} = \frac{m_q}{\varepsilon_p} \frac{m_q}{\varepsilon_{p'}} \mathcal{M}_{ps,p's'}^{ex} \quad (20)$$

Here the spin may be either parallel ($s = s'$) or anti-parallel ($s = -s'$). Thus scattering possibilities are denoted by $(+, +)$, $(+, -)$, $(-, -)$ etc. Motivated by [15], in analogy with isospin we define spin dependent interaction parameter as $f_{pp'}^+ = \frac{1}{2}(f_{pp'}^{++} + f_{pp'}^{+-})$ and $f_{pp'}^- = \frac{1}{2}(f_{pp'}^{--} + f_{pp'}^{-+})$. Note that, $f_{pp'}^{+-} = f_{pp'}^{-+}$.

For $(+, +)$ scattering the interaction parameter is given by

$$\begin{aligned} f_{pp'|p=p'=p_f^+}^{++} = & -\frac{g^2}{9\varepsilon_f^{+2}} \frac{1}{p_f^{+2}(1 - \cos \theta)} [2m_q^2 - p_f^{+2}(1 - \cos \theta) - p_f^{+2} \cos \theta_1 \cos \theta_2 \\ & + \frac{1}{(\varepsilon_f^+ + m_q)^2} \{m_q(\varepsilon_f^+ + m_q)p_f^{+2}(\cos^2 \theta_1 + \cos^2 \theta_2) + p_f^{+4} \cos \theta \cos \theta_1 \cos \theta_2\}], \end{aligned} \quad (21)$$

where $\hat{p} \cdot \hat{s} = \cos \theta_1$; $\hat{p}' \cdot \hat{s} = \cos \theta_2$ and Fermi energy $\varepsilon_f^+ = (p_f^{+2} + m_q^2)^{1/2}$. Since spin and momentum have no preferred direction, we have done angular average of the spin dependent parameter [24]:

$$\begin{aligned} \overline{f_{pp'|p=p'=p_f^+}^{++}} = & \int \frac{d\Omega_1}{4\pi} \int \frac{d\Omega_2}{4\pi} f_{pp'|p=p'=p_f^+}^{++} \\ = & -\frac{g^2}{9\varepsilon_f^{+2}p_f^{+2}(1 - \cos \theta)} \left[2m_q^2 - p_f^{+2}(1 - \cos \theta) + \frac{2m_q p_f^{+2}}{3(\varepsilon_f^+ + m_q)} \right]. \end{aligned} \quad (22)$$

¹ With the help of Eq.(5) along with the Eq.(22) one can find LPs, but it is to be noted that $f_{0,1}^{++}$ or $f_{0,1}^{--}$ are individually divergent because of the term, $(1 - \cos \theta)$, in the denominator of the interaction parameter. This divergence disappear if one uses Debye screening mass for

[1] denoted hereafter $\overline{f_{pp'}} = f_{pp'}$.

gluons or equivalently use HDL corrected gluon propagator while evaluating the scattering amplitudes [12, 24]. Note that the combination $\left(f_0^{++(--)}-\frac{1}{3}f_1^{++(--)}\right)$ is, however, finite as in this case the divergences cancel and we do not calculate the LPs separately. It would, however, be interesting to see how do the results modify if HDL calculations are performed to evaluate $f_{0,1}^{++(--)}$, $f_{0,1}^{+-}$ and the corresponding physical quantities. The numerical estimates suggest that for the results what we present here, the effect of HDL corrections are expected to be small.

From Eq.(5),

$$\begin{aligned} f_0^{++}-\frac{1}{3}f_1^{++} &= -\frac{g^2}{18\varepsilon_f^{+2}p_f^{+2}}\int_{-1}^{+1}\left[2m_q^2-p_f^{+2}(1-\cos\theta)+\frac{2m_qp_f^{+2}}{3(\varepsilon_f^++m_q)}\right]d(\cos\theta) \\ &= -\frac{g^2}{9\varepsilon_f^{+2}p_f^{+2}}\left[2m_q^2-p_f^{+2}+\frac{2m_qp_f^{+2}}{3(\varepsilon_f^++m_q)}\right]. \end{aligned} \quad (23)$$

The above combination will appear in the calculation of the chemical potential and other relevant quantities. For $(+, -)$ scattering, the angular averaged interaction parameter yields

$$f^{+-}\Big|_{p=p_f^+, p'=p_f^-} = \frac{g^2}{9\varepsilon_f^+\varepsilon_f^-}\left[1-\left\{\frac{m_qp_f^{+2}}{3(\varepsilon_f^++m_q)}+\frac{m_qp_f^{-2}}{3(\varepsilon_f^-+m_q)}\right\}\times\frac{1}{(m_q^2-\varepsilon_f^+\varepsilon_f^-+p_f^+p_f^-\cos\theta)}\right]. \quad (24)$$

It is to be noted that, individual LPs for scattering of unlike spin states are finite *i.e.* free of divergences, in contrast to the case involving scattering of like spin states (For details see Appendix).

A. Chemical potential

Now we proceed to calculate chemical potential, which, in principle, will be different for spin-up and spin-down quarks, denoted by μ^s with s (*or* s') = $+, -$ for matter containing unequal densities of up and down quarks. To determine the chemical potential with arbitrary polarization ξ , we take the distribution function with explicit spin index (s *or* s'), so that variation of distribution function gives [13, 20, 25]

$$\delta n_q^s = -N^s(0)\left[\sum_{s'}f_0^{ss'}\delta n_q^{s'}-\delta\mu^s\right], \quad (25)$$

where $N^s(0)$ is given by the Eq.(7). The Eq.(25) yields

$$\frac{\partial \mu^s}{\partial n_q^s} = \frac{1}{N^s(0)} + \sum_{s'} f_0^{ss'} \frac{\partial n_q^{s'}}{\partial n_q^s}. \quad (26)$$

Separately for spin-up and spin-down states we have

$$\begin{pmatrix} \partial \mu^+ \\ \partial \mu^- \end{pmatrix} = \begin{pmatrix} \frac{1}{N^+(0)} + f_0^{++} & f_0^{+-} \\ f_0^{-+} & \frac{1}{N^-(0)} + f_0^{--} \end{pmatrix} \begin{pmatrix} \partial n_q^+ \\ \partial n_q^- \end{pmatrix}, \quad (27)$$

where the superscripts $++$ and $+-$ denote scattering of quasiparticle with up-up and up-down spin states. For unpolarized matter the upper and lower component become equal which gives rise to the well known result [14]

$$\mu d\mu = \left[p_f + \frac{g_{deg} \mu p_f^2}{2\pi^2} (f_0 - \frac{1}{3} f_1) \right] dp_f. \quad (28)$$

In general the chemical potential (both for spin-up and spin-down) is the combination of like and unlike spin states. By adjusting the constant of integration [14], the chemical potential of spin-up quark turns out to be

$$\begin{aligned} \mu^+ = & \varepsilon_f^+ - \frac{g^2}{6\pi^2 \varepsilon_f^+} \left[\frac{11}{6} m_q^2 \ln \left(\frac{p_f^+ + \varepsilon_f^+}{m_q} \right) + \frac{2}{3} p_f^+ m_q - \frac{p_f^+ \varepsilon_f^+}{2} \right] \\ & + \frac{g^2}{72\pi^2 \varepsilon_f^+} \left[-\frac{2m_q^3}{p_f^+} \ln \left(\frac{p_f^+ + p_f^-}{p_f^+ - p_f^-} \right) + \frac{4m_q^2 \varepsilon_f^+}{p_f^+} \left\{ \ln \left(\frac{p_f^+ + p_f^-}{p_f^+ - p_f^-} \right) + \ln \left(\frac{p_f^+ \varepsilon_f^- + p_f^- \varepsilon_f^+}{p_f^+ \varepsilon_f^- - p_f^- \varepsilon_f^+} \right) \right\} \right. \\ & - 14m_q^2 \ln \left(\frac{p_f^- + \varepsilon_f^-}{m_q} \right) + 2m_q p_f^- - 3m_q p_f^- \ln \left(\frac{p_f^+ + p_f^-}{p_f^+ - p_f^-} \right) \\ & - \frac{m_q}{p_f^+} (2m_q^2 + 3p_f^{+2}) \ln \left(\frac{p_f^+ \varepsilon_f^- + p_f^- \varepsilon_f^+}{p_f^+ \varepsilon_f^- - p_f^- \varepsilon_f^+} \right) + 6m_q \varepsilon_f^+ \ln \left(\frac{p_f^- + \varepsilon_f^-}{m_q} \right) \\ & \left. + \frac{m_q}{p_f^+} \{ 2\varepsilon_f^- (2m_q - \varepsilon_f^+) - p_f^{-2} \} \ln \left(\frac{\varepsilon_f^- \varepsilon_f^+ - m_q^2 - p_f^- p_f^+}{\varepsilon_f^- \varepsilon_f^+ - m_q^2 + p_f^- p_f^+} \right) + 6p_f^- \varepsilon_f^- \right]. \quad (29) \end{aligned}$$

In the above equation the term in the first square bracket arises due to the scattering of like spin states $(++)$, while the latter comes from the scattering of unlike spin states $(+-)$.

Similarly, for spin-down quark, one may determine μ^- by replacing $p_f^\pm$ with $p_f^\mp$ and $\varepsilon_f^\pm$ with $\varepsilon_f^\mp$ in Eq.(29).

For the numerical estimation of the above mentioned quantities, following ref.[6, 26], we take $\alpha_c = g^2/4\pi = 2.2$, is the fine structure constant of QCD and $m_q = 300\text{MeV}$. In Fig(1) we plot chemical potential for spin-up and spin-down quark as a function of density with order parameter $\xi = 0.5$. In real astrophysical calculations, the chemical potentials are determined by the β -equilibrium conditions where the condition of charge neutrality is also imposed. In Fig(1), we, however, use density n_q and polarization parameter ξ as input parameters and Eq.(29) is used to determine μ for a system with one flavor.




FIG. 1: Density dependence of chemical potential of spin-up and spin-down quark denoted by solid and dashed curve respectively.

B. Energy density

Once the μ is determined, one can readily calculate the exchange energy density by evaluating [14, 17, 25]

$$E_{ex} = \int dn_q (\mu - \varepsilon_f) \quad (30)$$

After summing up over the color degrees of freedom and evaluating over the Fermi surfaces, we have the exchange energy density. The latter consisting of all types of scattering

amplitudes, can be written as

$$E_{ex} = E_{ex}^{++} + E_{ex}^{--} + E_{ex}^{+-}, \quad (31)$$

which we evaluate numerically. The total kinetic energy density for spin-up and spin-down quark is given by

$$E_{kin} = \frac{3}{16\pi^2} \sum_{s=\pm} \left[p_f^s \varepsilon_f^s (\varepsilon_f^{s^2} + p_f^{s^2}) - m_q^4 \ln \left(\frac{\varepsilon_f^s + p_f^s}{m_q} \right) \right], \quad (32)$$

where $\varepsilon_f^s = (p_f^{s^2} + m_q^2)^{1/2}$. The total energy is given by the sum of the kinetic energy and the interaction energy E_{ex} *i.e.*

$$E_{tot} = E_{kin} + E_{ex}. \quad (33)$$

Now we calculate incompressibility and sound velocity by using Eq.(12) and Eq.(14). In Fig.(2) and Fig.(3) we plot the density dependence of incompressibility and sound velocity. This shows for higher value of the order parameter ξ , the incompressibility and the sound velocity becomes higher for the same value of density. Thus the EOS for polarized quark matter is found to be stiffer than the unpolarized one.




FIG. 2: Incompressibility K in quark matter as a function of density for different polarization parameter.




FIG. 3: First sound velocity c_1 in quark matter as a function of density for different polarization parameter.

C. Phase transition

Bloch first pointed out the possibility of ferromagnetism of electron gas where the Fock exchange interaction induces spontaneous spin polarization [27]. Consider the spin polarized electron gas interacting by the Coulomb interaction in the background of the positively charged ions. Since the direct interaction gives no contribution due to charge neutrality, the Fock exchange interaction gives the leading contribution as the interaction energy. For spontaneous ferromagnetism, the interaction energy dominates over the kinetic energy [6, 21, 28].

Therefore, if the exchange energy due to OGE interaction is negative and becomes greater than the kinetic energy at some density, the quark matter becomes polarized giving rise to ferromagnetism [6].

To check whether our results for the total energy density are consistent with ref.[6], we consider two limiting cases corresponding to the ultra-relativistic (UR) and non-relativistic (NR) regimes. In the ultra-relativistic (UR) limit, $p_f^s \gg m_q$, then using Eq.(29) we have

$$\mu^{+,ur} = p_f^+ + \frac{\alpha_c}{3\pi} \left[p_f^+ + \frac{p_f^{-2}}{p_f^+} \right], \quad (34)$$

Similarly one can find $\mu^{-,ur}$ by replacing $p_f^\pm$ with $p_f^\mp$.

One can arrive at the same expression $\mu^{\pm,ur}$ by taking UR limit of the scattering amplitude. For $(+, +)$ scattering one gets the interaction parameter as

$$f_{pp'}^{++,ur} = \frac{g^2}{9pp'} (1 + \cos \theta_1 \cos \theta_2). \quad (35)$$

After taking angular average of the interaction parameter and with the help of Eq.(5), we find that f_1^{++} vanishes. Thus we have

$$f_{pp'}^{++,ur} \Big|_{p=p'=p_f^+} = f_0^{++,ur} = \frac{g^2}{9p_f^{+2}}. \quad (36)$$

Similarly for $(+, -)$ scattering, the interaction parameter yields

$$f_{pp'}^{+-,ur} = \frac{g^2}{9pp'} (1 - \cos \theta_1 \cos \theta_2). \quad (37)$$

The only existing LP is f_0^{+-} and other higher order LPs does not contribute. Hence we get

$$f_{pp'}^{+-,ur} \Big|_{p=p_f^+, p'=p_f^-} = f_0^{+-,ur} = \frac{g^2}{9p_f^+ p_f^-}. \quad (38)$$

It is observed that, in UR limit, all the LPs are finite. Now the chemical potential for spin-up quark is found to be

$$\mu^{+,ur} = p_f^+ + \frac{\alpha_c}{3\pi} \left[p_f^+ + \frac{p_f^{-2}}{p_f^+} \right]. \quad (39)$$

The chemical potential, $\mu^{-,ur}$, can be obtained by replacing $p_f^{\pm}$ with $p_f^{\mp}$ in Eq.(39).

Using Eqs.(30) and (31), the exchange energy densities are given by

$$\left. \begin{aligned} E_{ex}^{++,ur} &= \frac{\alpha_c}{8\pi^3} p_f^{+4} \\ E_{ex}^{--,ur} &= \frac{\alpha_c}{8\pi^3} p_f^{-4} \\ E_{ex}^{+-,ur} &= \frac{\alpha_c}{4\pi^3} p_f^{+2} p_f^{-2} \end{aligned} \right\} \quad (40)$$

Thus the final expression for the exchange energy density in the UR limit is found to be

$$E_{ex}^{ur} = \frac{\alpha_c}{8\pi^3} p_f^4 [(1 + \xi)^{4/3} + (1 - \xi)^{4/3} + 2(1 - \xi^2)^{2/3}]. \quad (41)$$

This result is same as in ref.[6].

Similarly from Eq.(32), the kinetic energy density in UR limit takes the following form [6]:

$$E_{kin}^{ur} = \frac{3p_f^4}{8\pi^2} [(1 + \xi)^{4/3} + (1 - \xi)^{4/3}]. \quad (42)$$

In the NR limit, $p(\text{or } p') \ll m_q$, the interaction parameter reduces to a simple form,

$$f_{ps,p's'}^{nr} = -\frac{g^2}{9pp'} \left[\frac{1 + s \cdot s'}{(1 - \cos \theta)} \right]. \quad (43)$$

For spin anti-parallel interaction $s = -s'$, then $f_{ps,p's'}^{nr} = 0$. Thus the contribution due to the scattering of quarks with unlike spin states vanishes and the dominant contribution to energy density comes from the parallel spin states ($s = s'$). For the (s, s) scattering, the interaction parameter yields

$$f_{pp'}^{nr,s} \Big|_{p=p'=p_f^s} = -\frac{2g^2}{9p_f^{s^2}(1 - \cos \theta)}, \quad (44)$$

where $s = +$ or $-$ according to scattering process. In NR limit one gets

$$(f_0^s - \frac{1}{3}f_1^s) = -\frac{2g^2}{9p_f^{s^2}}. \quad (45)$$

The NR chemical potential μ^{nr} is given by

$$\mu^{nr,s} = m_q - \frac{g^2}{3\pi^2} p_f^s. \quad (46)$$

Using Eq.(30), the exchange energy density for the $(+, +)$ scattering is given by

$$E_{ex}^{++,nr} = -\frac{g^2}{8\pi^4} p_f^4 (1 + \xi)^{4/3}. \quad (47)$$

Similarly for $(-, -)$ scattering, we have

$$E_{ex}^{--,nr} = -\frac{g^2}{8\pi^4} p_f^4 (1 - \xi)^{4/3}. \quad (48)$$

As in the NR limit $E_{ex}^{+-} = E_{ex}^{-+} = 0$ as mentioned before, so from Eq.(31) the exchange energy density yields

$$E_{ex}^{nr} = -\frac{\alpha_c}{2\pi^3} p_f^4 [(1 + \xi)^{4/3} + (1 - \xi)^{4/3}]. \quad (49)$$

Thus the energy density, in this limit, becomes negative. The kinetic energy density turns out to be [6]

$$E_{kin}^{nr} = \frac{3p_f^5}{20\pi^2 m_q} [(1 + \xi)^{5/3} + (1 - \xi)^{5/3}]. \quad (50)$$

In NR limit ferromagnetism can appear as a consequence of competition between the kinetic energy and the Coulomb potential energy [10]. The latter favors the spin alignment due

to quantum effect. When the energy gain due to the spin alignment dominate over the increase in the kinetic energy at some density, the unpolarized state suddenly turns into the completely polarized state [29]. On the other hand, in UR limit, the contribution to the energy density not only comes from the like spin states but also unlike spin states of scatterer (see [6] for detailed discussion).




FIG. 4: The total energy of quark liquid as a function of polarization parameter at $n_q = 0.1 \text{ fm}^{-3}$, $n_q = 0.14 \text{ fm}^{-3}$ and $n_q = 0.2 \text{ fm}^{-3}$. The critical density is found to be $n_q^c = 0.14 \text{ fm}^{-3}$ in this case.

To check the consistency we compare our result derived in RFLT approach with that of ref[6] derived from two loop ring diagrams. In Fig.(4) we plot E_{tot}/n_q as a function of polarization parameter ξ . The results clearly show that for lower density ($< 0.14 \text{ fm}^{-3}$), total energy favors at $\xi = 1$ which indicates completely polarized state; while at higher density, the system becomes unpolarized ($\xi = 0$). Thus the polarization parameter suddenly changes from $\xi = 1$ to $\xi = 0$ as one increases the number density of the system. So the phase transition is first order and the critical density n_q^c is around 0.14 fm^{-3} .

In Fig.(5) we show total energy as a function of polarization parameter for different densities. In every plot, there is a minima which corresponds to a possible metastable state. We notice that when density increases metastable state arises for lower values of polarization parameter ξ . For example, at density $\sim 0.2 \text{ fm}^{-3}$ minima arises at $\xi = 0.1$ while at density $\sim 0.35 \text{ fm}^{-3}$ minima arises at $\xi = 0.03$. Thus the metastable state shows a tendency of




FIG. 5: Metastable ferromagnetic state as a function of polarization parameter for different densities.

disappearance as the density increased.

IV. SUMMARY AND CONCLUSION

To summarize and conclude, in this work we have applied RFLT to study the properties of dense quark matter. Accordingly, we calculate the FLPs by retaining their explicit spin dependencies. We also show how the physical quantities like chemical potential of spin up and spin down states, their energy densities and the quantities like incompressibility, sound velocity for polarized quark matter can be expressed in terms of these spin dependent RFLPs. For the scattering involving like spin states, the LPs $f_{0,1}^{++}$ and $f_{0,1}^{--}$ are found to

diverge. However, we show that the combination in which they appear in the calculation of the physical quantities such divergences cancel. For the scattering involving unlike spin states no such divergence appear. The appearance of such divergences is related to the unscreened gluonic interaction between the quarks which might be cured by invoking hard dense loop corrected gluon propagator. We do not perform such calculation here and postpone this for future investigation. As far as the equation of state (EOS) is concerned, we in the present model find that the EOS for the polarized quark matter is stiffer than the unpolarized one. In addition, we also show that there exists a metastable state which disappear at higher density, although it seems that the effect is tiny.

We reconfirm that DQM can exhibit ferromagnetism at low density as was originally suggested in [6]. However, the density at which the spin polarized ferromagnetic state in the present model might appear depends strongly on the quark mass. The critical density increases with increasing mass. In Fig. (4), we observe that states with ξ appear only below or around normal nuclear density where deconfined quark matter is not likely to exist. We cannot, however, ascertain the critical density from the present analysis where we restrict ourselves only to OGE diagrams and one flavor system. In this regime, multi-gluon exchange processes [10] might play an important role. Furthermore, the correlations as given by the ring diagrams can also change the conclusion. Further work therefore is necessary to understand the existence of ferromagnetic quark matter in real multi flavor system which might appear in astrophysics.

V. APPENDIX

In the text the interaction parameter $f_{pp'}^{+-}$ for unlike spin states was calculated. Here we give detail expression of Landau parameters. With the help of Eq.(5), the LPs are given by,

$$f_0^{+-} = \frac{g^2}{18\varepsilon_f^+\varepsilon_f^-} \left[2 + \frac{m_q[\varepsilon_f^-p_f^{+2} + m_q(p_f^{+2} + p_f^{-2}) + \varepsilon_f^+p_f^{-2}]}{3p_f^+p_f^-(m_q + \varepsilon_f^+)(m_q + \varepsilon_f^-)} \ln \left(\frac{m_q^2 - p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-}{m_q^2 + p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-} \right) \right] \quad (51)$$

and

$$f_1^{+-} = \frac{g^2}{18\varepsilon_f^+\varepsilon_f^-} \left[6 - \frac{2m_q[\varepsilon_f^-p_f^{+2} + m_q(p_f^{+2} + p_f^{-2}) + \varepsilon_f^+p_f^{-2}]}{p_f^+p_f^-(m_q + \varepsilon_f^+)(m_q + \varepsilon_f^-)} + \right. \\ \left. \left(\frac{m_q(m_q^2 - \varepsilon_f^+\varepsilon_f^-)[\varepsilon_f^-p_f^{+2} + m_q(p_f^{+2} + p_f^{-2}) + \varepsilon_f^+p_f^{-2}]}{p_f^{+2}p_f^{-2}(m_q + \varepsilon_f^+)(m_q + \varepsilon_f^-)} \right) \ln \left(\frac{m_q^2 + p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-}{m_q^2 - p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-} \right) \right]. \quad (52)$$

Using Eq.(51) and Eq.(52) we have

$$f_0^{+-} - \frac{1}{3}f_1^{+-} = \frac{g^2}{18\varepsilon_f^+\varepsilon_f^-} \left[2 - \left\{ \frac{m_qp_f^{+2}}{3(\varepsilon_f^+ + m_q)} + \frac{m_qp_f^{-2}}{3(\varepsilon_f^- + m_q)} \right\} \times \right. \\ \left. \left\{ -\frac{2}{p_f^+p_f^-} + \frac{(p_f^+p_f^- + m_q^2 - \varepsilon_f^+\varepsilon_f^-)}{(p_f^{+2}p_f^{-2})} \ln \left(\frac{m_q^2 + p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-}{m_q^2 - p_f^+p_f^- - \varepsilon_f^+\varepsilon_f^-} \right) \right\} \right]. \quad (53)$$

Acknowledgments

The authors would like to thank P.Roy for his critical reading of the manuscript.

-
- [1] R.C.Hwa, *Quark Gluon Plasma-Vol-1*, (World Scientific) (1990).
R.C.Hwa, *Quark Gluon Plasma-Vol-2*, (World Scientific) (1995).
R.C.Hwa et al., *Quark Gluon Plasma-Vol-3*, (World Scientific) (2004).
 - [2] A.Niegawa, Prog.Theor.Phys **113**, 581 (2005).
 - [3] M.G.Alford, A.Schmitt, K.Rajagopal and T.Schaffer, arXiv: 0709.4635 [hep-ph], (2007).
M.G.Alford, J.A.Bowers and K.Rajagopal, J.Phys.**G27**, 541 (2001).
 - [4] M.Inui, H.Kohyama and A.Niegawa, arXiv: 0709.2204 [hep-ph], (2007).
 - [5] E.Nakano, T.Maruyama and T.Tatsumi, Phys.Rev.**D68**, 105001 (2003).
 - [6] T.Tatsumi, Phys.Lett.**B 489**, 280 (2000).
 - [7] T.Maruyama and T.Tatsumi, Nucl.Phys.**A693**, 710 (2001).
 - [8] D.T.Son and M.A.Stephanov, Phys.Rev.**D77**, 014021 (2008).
 - [9] T.Tatsumi, T.Maruyama, E.Nakano and K.Nawa, Nucl.Phys.**A 774**, 827 (2006).
 - [10] K.Ohnishi, M.Oka and S.Yasui, Phys.Rev.**D76**, 097501 (2007).

- [11] S.Hands, S.Kim, J.I.Skullerud, Eur.Phys.J **C48**, 193 (2006).
- [12] T.Tatsumi and K.Sato, Phys.Lett **B663**, 322 (2008).
- [13] G.Baym and C.Pethick, *Landau Fermi liquid Theory: Concepts and Applications*, United States Of America (1991).
- [14] G.Baym and S.A.Chin, Nucl.Phys.**A262**, 527 (1976).
- [15] A.B.Migdal, *Theory of Finite Fermi Systems* (1967) (Wiley, New York) (Russ.ed.1965),
A.B.Migdal, *Nuclear Theory: The Quasiparticle Method* (1968) (Benjamin, New York).
- [16] T.Matsui, Nucl.Phys.**A370**, 365 (1981).
- [17] S.A.Chin, Annals of Physics **108**, 301 (1977).
- [18] A.Schwenk, G.E.brown and B.Friman, Nucl.Phys. **A703**, 745 (2002).
- [19] J.W.Negele and H.Orland, *Quantum many particle systems*, Addison-Wesley Publishing Company, (1988).
- [20] R.M.Aguirre and A.L.De.Paoli, Phys.Rev.**C75**, 045207 (2007).
- [21] T.Tatsumi, PoS,031 (JHW 2005).
- [22] V.B.Berestetsii, E.M.Lifshitz and L.P.Pitaevsii, *Relativistic Quantum Theory*, Pergamon Press, (1971).
- [23] Elliot Leader, *Introduction to gauge theories and modern particle physics-Vol-1 and Vol-2*, (Cambridge University Press), (1996).
- [24] T.Tatsumi, arXiv:0711.3349v1[hep-ph], (2007).
- [25] K.Pal, S.Biswas and A.K. Dutt-Mazumder, arXiv: 0803.3392 [hep-ph], (2008).
- [26] T. DeGrand, R. L. Jaffe, K. Johnson and J. Kiskis, Phys. Rev. **D12**, 2060 (1975).
- [27] F.Bloch, Z.Phys.**57**, 545 (1929).
- [28] T.Tatsumi, arXiv: nucl-th/000201v1, (2000).
- [29] T.Tatsumi, arXiv: astro-ph/0004062v1, (2000).